

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

What Is Claimed Is:


1. A compound of Formula I:



wherein each of R^1 and R^{11} is a group independently selected from hydrido, alkyl, alkylaminoalkyl and phenyl; wherein p is a number selected from zero through five, inclusive; wherein r is a number selected from zero, one and two; wherein R^2 is selected from hydrido and alkyl; wherein R^3 is a group selected from hydrido, cycloalkylalkyl, aralkyl and haloaralkyl; wherein each of R^4 and R^6 is a group independently selected from hydrido and methyl; wherein R^5 is selected from

10

15




wherein V is selected from hydrido, alkyl, cycloalkyl, aryl and aralkyl; wherein each of R^9 and R^{10} is a group independently selected from hydrido, alkyl, alkenyl, alkynyl, cycloalkyl and aryl; wherein m is a number selected from zero through three; wherein n is a number selected from zero through three; wherein R^7 is a group selected from alkyl, cycloalkylalkyl and aralkyl; wherein R^8 is a group selected from hydrido, alkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, alkenyl and haloalkenyl; wherein each of R^{12} and R^{13} is a group independently selected from hydrido, alkyl, cycloalkyl,

20

25

cycloalkylalkyl, alkylacyl, aryl, aralkyl, haloaryl and haloaralkyl; and wherein any one of said R¹ through R¹³ groups having a substitutable position may be substituted with one or more groups selected from alkyl, hydroxy, hydroxyalkyl, halo, alkoxy, alkoxyalkyl and alkenyl; or a pharmaceutically-acceptable salt thereof.


2. Compound of Claim 1 wherein each of R¹ and R¹¹ is independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, N,N'-dimethylaminomethyl, N,N'-diethylaminomethyl, N,N'-diethylaminoethyl and phenyl; wherein p is a number selected from zero through four, inclusive; wherein r is a number selected from zero, one and two; wherein R² is selected from hydrido and alkyl; wherein R³ is selected from hydrido, cycloalkylalkyl, phenylalkyl, halophenylalkyl, naphthylalkyl and halonaphthylalkyl; wherein each of R⁴ and R⁶ is independently selected from hydrido and methyl; wherein R⁵ is selected from



wherein V is selected from hydrido, alkyl, phenyl and benzyl; wherein each of R⁹ and R¹⁰ is independently selected from hydrido, alkyl, alkenyl, alkynyl, cycloalkyl and aryl; wherein m is a number selected from zero through three; wherein n is a number selected from zero through three; wherein R⁷ is selected from cyclohexylmethyl and benzyl, either one of which may be substituted with one or more groups selected from alkyl, hydroxy and alkoxy; wherein R⁸ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkenyl

and haloalkenyl; and wherein each of R^{12} and R^{13} is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkanoyl, halophenyl, phenylalkyl, halophenylalkyl, naphthyl, halonaphthyl, naphthylalkyl and halonaphthylalkyl; or a pharmaceutically-acceptable salt thereof.


3. Compound of Claim 2 wherein each of R^1 and R^{11} is independently selected from hydrido, methyl, ethyl, n-propyl and isopropyl; wherein p is a number selected from zero through three, inclusive; wherein r is a number selected from zero, one and two; wherein R^2 is selected from hydrido, methyl, ethyl and n-propyl; wherein R^3 is selected from hydrido, cyclohexylmethyl, benzyl, phenylethyl, fluorobenzyl, fluorophenylethyl, chlorobenzyl, chlorophenylethyl, naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and chloronaphthylmethyl; wherein each of R^4 and R^6 is independently selected from hydrido and methyl; wherein R^5 is selected from



wherein V is selected from hydrido, alkyl, cycloalkyl, aryl and aralkyl; wherein m is a number selected from one through three; wherein R^7 is cyclohexylmethyl; wherein R^8 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, cyclobutyl, cyclopropylmethyl, cyclobutylmethyl, cyclohexylmethyl, allyl and vinyl; and wherein each of R^{12} and R^{13} is independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl,


tert-butyl, cyclopropyl, cyclopropylmethyl,
 cyclopropylethyl, propylcarbonyl, ethylcarbonyl,
 methylcarbonyl, phenyl, benzyl, phenylethyl,
 monochlorophenyl, dichlorophenyl, monofluorophenyl,
 5 difluorophenyl, monochlorophenylmethyl,
 monochlorophenylethyl, dichlorophenylmethyl,
 dichlorophenylethyl, naphthyl, monofluoronaphthyl,
 monochloronaphthyl, naphthylmethyl, naphthylethyl,
 fluoronaphthylmethyl and chloronaphthylethyl; or a
 10 pharmaceutically-acceptable salt thereof.

4. Compound of Claim 3 wherein each of R^1 and
 R^{11} is independently hydrido or methyl; wherein p is a
 number selected from zero through three, inclusive;
 15 wherein r is zero or two; wherein R^2 is selected from
 hydrido, methyl, ethyl and n-propyl; wherein R^3 is
 selected from hydrido, cyclohexylmethyl, benzyl,
 phenylethyl, phenylpropyl, fluorobenzyl,
 fluorophenylethyl, chlorobenzyl, chlorophenylethyl,
 20 naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and
 chloronaphthylmethyl; wherein each of R^4 and R^6 is
 hydrido; wherein R^5 is selected from




25 wherein V is selected from hydrido and methyl; wherein m
 is one or two; wherein R^7 is cyclohexylmethyl; wherein R^8
 is selected from ethyl, n-propyl, n-butyl, isobutyl,
 cyclopropyl, cyclobutyl, cyclopropylmethyl, allyl and
 30 vinyl; wherein each of R^{12} and R^{13} is independently
 selected from hydrido, methyl, ethyl, n-propyl,
 isopropyl, cyclopropylmethyl, phenyl, benzyl,
 monochlorophenyl and dichlorophenyl; or a
 pharmaceutically-acceptable salt thereof.

5. Compound of Claim 4 of Formula II




5 wherein r is zero or two; wherein q is two or three;
 wherein R^2 is selected from hydrido, methyl, ethyl and
 phenyl; wherein R^3 is selected from hydrido,
 cyclohexylmethyl, benzyl, fluorobenzyl, chlorobenzyl,
 fluoronaphthylmethyl and chloronaphthylmethyl; wherein
 10 each of R^4 and R^6 is hydrido; wherein R^5 is selected from







wherein V is selected from hydrido and methyl; wherein m is one or
 15 two; wherein R^7 is cyclohexylmethyl; wherein R^8 is
 selected from n -propyl, isobutyl, cyclopropyl,
 cyclopropylmethyl, allyl and vinyl; wherein R^{12} and R^{13} is
 independently selected from methyl, ethyl and isopropyl;
 or a pharmaceutically-acceptable salt thereof.





20


6. Compound of Claim 5 selected from compounds,
 their tautomers and pharmaceutically-acceptable salts
 thereof, of the group consisting of:



25








5

7. Compound of Claim 6 which is N-[1R*-
[[[1S,1R*-(cyclohexylmethyl)-2S*,3R*-dihydroxy-5-


methylhexyl]amino]carbonyl]-3-butynyl]- α R*-[[[2-(dimethylamino)ethyl]sulfonyl]methyl]benzenepropanamide or a pharmaceutically-acceptable salt thereof.

- 5 8. A pharmaceutical composition comprising a therapeutically-effective amount of a renin-inhibiting compound and a pharmaceutically-acceptable carrier or diluent, said renin-inhibiting compound selected from a family of compounds of Formula I:

10




- wherein each of R^1 and R^{11} is a group independently selected from hydrido, alkyl, alkylaminoalkyl and phenyl;
 15 wherein p is a number selected from zero through five, inclusive; wherein r is a number selected from zero, one and two; wherein R^2 is selected from hydrido and alkyl; wherein R^3 is a group selected from hydrido, cycloalkylalkyl, aralkyl and haloaralkyl; wherein each of
 20 R^4 and R^6 is a group independently selected from hydrido and methyl; wherein R^5 is selected from



- 25 wherein V is selected from hydrido, alkyl, cycloalkyl, aryl and aralkyl; wherein each of R^9 and R^{10} is a group independently selected from hydrido, alkyl, alkenyl, alkynyl, cycloalkyl and aryl; wherein m is a number selected from zero through three; wherein n is a number

selected from zero through three; wherein R^7 is a group selected from alkyl, cycloalkylalkyl and aralkyl; wherein R^8 is a group selected from hydrido, alkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, alkenyl and haloalkenyl;
 5 wherein each of R^{12} and R^{13} is a group independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkylacyl, aryl, aralkyl, haloaryl and haloaralkyl; and wherein any one of said R^1 through R^{13} groups having a substitutable position may be substituted
 10 with one or more groups selected from alkyl, hydroxy, hydroxyalkyl, halo, alkoxy, alkoxyalkyl and alkenyl; or a pharmaceutically-acceptable salt thereof.


9. The composition of Claim 8 wherein each of
 15 R^1 and R^{11} is independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, N,N'-dimethylaminomethyl, N,N'-diethylaminomethyl, N,N'-diethylaminoethyl and phenyl; wherein p is a number selected from zero through four,
 20 inclusive; wherein r is a number selected from zero, one and two; wherein R^2 is selected from hydrido and alkyl; wherein R^3 is selected from hydrido, cycloalkylalkyl, phenylalkyl, halophenylalkyl, naphthylalkyl and halonaphthylalkyl; wherein each of R^4 and R^6 is
 25 independently selected from hydrido and methyl; wherein R^5 is selected from



30 wherein V is selected from hydrido, alkyl, phenyl and benzyl; wherein each of R^9 and R^{10} is independently selected from hydrido, alkyl, alkenyl, alkynyl,

cycloalkyl and aryl; wherein m is a number selected from zero through three; wherein n is a number selected from zero through three; wherein R⁷ is selected from cyclohexylmethyl and benzyl, either one of which may be substituted with one or more groups selected from alkyl, hydroxy and alkoxy; wherein R⁸ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkenyl and haloalkenyl; and wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkanoyl, halophenyl, phenylalkyl, halophenylalkyl, naphthyl, halonaphthyl, naphthylalkyl and halonaphthylalkyl; or a pharmaceutically-acceptable salt thereof.


10. The composition of Claim 9 wherein each of R¹ and R¹¹ is independently selected from hydrido, methyl, ethyl, n-propyl and isopropyl; wherein p is a number selected from zero through three, inclusive; wherein r is a number selected from zero, one and two; wherein R² is selected from hydrido, methyl, ethyl and n-propyl; wherein R³ is selected from hydrido, cyclohexylmethyl, benzyl, phenylethyl, fluorobenzyl, fluorophenylethyl, chlorobenzyl, chlorophenylethyl, naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and chloronaphthylmethyl; wherein each of R⁴ and R⁶ is independently selected from hydrido and methyl; wherein R⁵ is selected from



wherein V is selected from hydrido, alkyl, cycloalkyl, aryl and aralkyl; wherein m is a number selected from one

through three; wherein R^7 is cyclohexylmethyl; wherein R^8 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, cyclobutyl, cyclopropylmethyl, cyclobutylmethyl, cyclohexylmethyl, allyl and vinyl; and wherein each of R^{12} and R^{13} is independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, cyclopropylmethyl, cyclopropylethyl, propylcarbonyl, ethylcarbonyl, methylcarbonyl, phenyl, benzyl, phenylethyl, monochlorophenyl, dichlorophenyl, monofluorophenyl, difluorophenyl, monochlorophenylmethyl, monochlorophenylethyl, dichlorophenylmethyl, dichlorophenylethyl, naphthyl, monofluoronaphthyl, monochloronaphthyl, naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and chloronaphthylethyl; or a pharmaceutically-acceptable salt thereof.

11. The composition of Claim 10 wherein each of R^1 and R^{11} is independently hydrido or methyl; wherein p is a number selected from zero through three, inclusive; wherein r is zero or two; wherein R^2 is selected from hydrido, methyl, ethyl and n-propyl; wherein R^3 is selected from hydrido, cyclohexylmethyl, benzyl, phenylethyl, phenylpropyl, fluorobenzyl, fluorophenylethyl, chlorobenzyl, chlorophenylethyl, naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and chloronaphthylmethyl; wherein each of R^4 and R^6 is hydrido; wherein R^5 is selected from




wherein V is selected from hydrido and methyl; wherein m is one or two; wherein R^7 is cyclohexylmethyl; wherein R^8 is selected from ethyl, n-propyl, n-butyl, isobutyl, cyclopropyl, cyclobutyl, cyclopropylmethyl, allyl and vinyl; wherein each of R^{12} and R^{13} is independently

selected from hydrido, methyl, ethyl, n-propyl, isopropyl, cyclopropylmethyl, phenyl, benzyl, monochlorophenyl and dichlorophenyl; or a pharmaceutically-acceptable salt thereof.

5


12. The composition of Claim 11 wherein said compound is of Formula II



10

wherein r is zero or two; wherein q is two or three; wherein R² is selected from hydrido, methyl, ethyl and phenyl; wherein R³ is selected from hydrido, cyclohexylmethyl, benzyl, fluorobenzyl, chlorobenzyl, fluoronaphthylmethyl and chloronaphthylmethyl; wherein each of R⁴ and R⁶ is hydrido; wherein R⁵ is selected from

15



20


wherein V is selected from hydrido and methyl; wherein m is one or two; wherein R⁷ is cyclohexylmethyl; wherein R⁸ is


selected from n-propyl, isobutyl, cyclopropyl, cyclopropylmethyl, allyl and vinyl; wherein R^{12} and R^{13} is independently selected from methyl, ethyl and isopropyl; or a pharmaceutically-acceptable salt thereof.


5


13. The composition of Claim 12 wherein said renin inhibitor compound is selected from compounds, their tautomers and pharmaceutically-acceptable salts thereof, of the group consisting of:

10










14. The composition of Claim 13 wherein
 5 said renin-inhibitor compound is N-[1R*-[[[1S,1R*-(cyclohexylmethyl)-2S*,3R*-dihydroxy-5-methylhexyl]amino]carbonyl]-3-butynyl]-αR*-[[[2-dimethylamino)ethyl]sulfonyl]methyl]benzenepropanamide.


10

15. A therapeutic method for treating hypertension or glaucoma, said method comprising administering to a hypertensive patient a therapeutically-effective amount of a compound of Formula
 15 I:



wherein each of R¹ and R¹¹ is a group independently
 20 selected from hydrido, alkyl, alkylaminoalkyl and phenyl;

wherein p is a number selected from zero through five, inclusive; wherein r is a number selected from zero, one and two; wherein R² is selected from hydrido and alkyl; wherein R³ is a group selected from hydrido, cycloalkylalkyl, aralkyl and haloaralkyl; wherein each of R⁴ and R⁶ is a group independently selected from hydrido and methyl; wherein R⁵ is selected from




10

wherein V is selected from hydrido, alkyl, cycloalkyl, aryl and aralkyl; wherein each of R⁹ and R¹⁰ is a group independently selected from hydrido, alkyl, alkenyl, alkynyl, cycloalkyl and aryl; wherein m is a number selected from zero through three; wherein n is a number selected from zero through three; wherein R⁷ is a group selected from alkyl, cycloalkylalkyl and aralkyl; wherein R⁸ is a group selected from hydrido, alkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, alkenyl and haloalkenyl; wherein each of R¹² and R¹³ is a group independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkylacyl, aryl, aralkyl, haloaryl and haloaralkyl; and wherein any one of said R¹ through R¹³ groups having a substitutable position may be substituted with one or more groups selected from alkyl, hydroxy, hydroxyalkyl, halo, alkoxy, alkoxyalkyl and alkenyl; or a pharmaceutically-acceptable salt thereof.

16. The method of Claim 15 wherein each of R¹ and R¹¹ is independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, N,N'-dimethylaminomethyl, N,N'-diethylaminomethyl, N,N'-diethylaminoethyl and phenyl;

wherein p is a number selected from zero through four, inclusive; wherein r is a number selected from zero, one and two; wherein R² is selected from hydrido and alkyl; wherein R³ is selected from hydrido, cycloalkylalkyl, phenylalkyl, halophenylalkyl, naphthylalkyl and halonaphthylalkyl; wherein each of R⁴ and R⁶ is independently selected from hydrido and methyl; wherein R⁵ is selected from



10


wherein V is selected from hydrido, alkyl, phenyl and benzyl; wherein each of R⁹ and R¹⁰ is independently selected from hydrido, alkyl, alkenyl, alkynyl, cycloalkyl and aryl; wherein m is a number selected from zero through three; wherein n is a number selected from zero through three; wherein R⁷ is selected from cyclohexylmethyl and benzyl, either one of which may be substituted with one or more groups selected from alkyl, hydroxy and alkoxy; wherein R⁸ is selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkenyl and haloalkenyl; and wherein each of R¹² and R¹³ is independently selected from hydrido, alkyl, cycloalkyl, cycloalkylalkyl, alkanoyl, halophenyl, phenylalkyl, halophenylalkyl, naphthyl, halonaphthyl, naphthylalkyl and halonaphthylalkyl; or a pharmaceutically-acceptable salt thereof.

17. The method of Claim 16 wherein each of R¹ and R¹¹ is independently selected from hydrido, methyl, ethyl, n-propyl and isopropyl; wherein p is a number selected from zero through three, inclusive; wherein r is

30


a number selected from zero, one and two; wherein R^2 is selected from hydrido, methyl, ethyl and n-propyl; wherein R^3 is selected from hydrido, cyclohexylmethyl, benzyl, phenylethyl, fluorobenzyl, fluorophenylethyl, chlorobenzyl, chlorophenylethyl, naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and chloronaphthylmethyl; wherein each of R^4 and R^6 is independently selected from hydrido and methyl; wherein R^5 is selected from

10




wherein V is selected from hydrido, alkyl, cycloalkyl, aryl and aralkyl; wherein m is a number selected from one through three; wherein R^7 is cyclohexylmethyl; wherein R^8 is selected from methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, cyclobutyl, cyclopropylmethyl, cyclobutylmethyl, cyclohexylmethyl, allyl and vinyl; and wherein each of R^{12} and R^{13} is independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, cyclopropylmethyl, cyclopropylethyl, propylcarbonyl, ethylcarbonyl, methylcarbonyl, phenyl, benzyl, phenylethyl, monochlorophenyl, dichlorophenyl, monofluorophenyl, difluorophenyl, monochlorophenylmethyl, monochlorophenylethyl, dichlorophenylmethyl, dichlorophenylethyl, naphthyl, monofluoronaphthyl, monochloronaphthyl, naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and chloronaphthylethyl; or a pharmaceutically-acceptable salt thereof.

18. The method of Claim 17 wherein each of R^1 and R^{11} is independently hydrido or methyl; wherein p is a number selected from zero through three, inclusive; wherein r is zero or two; wherein R^2 is selected from hydrido, methyl, ethyl and n -propyl; wherein R^3 is selected from hydrido, cyclohexylmethyl, benzyl, phenylethyl, phenylpropyl, fluorobenzyl, fluorophenylethyl, chlorobenzyl, chlorophenylethyl, naphthylmethyl, naphthylethyl, fluoronaphthylmethyl and chloronaphthylmethyl; wherein each of R^4 and R^6 is hydrido; wherein R^5 is selected from




wherein V is selected from hydrido and methyl; wherein m is one or two; wherein R^7 is cyclohexylmethyl; wherein R^8 is selected from ethyl, n -propyl, n -butyl, isobutyl, cyclopropyl, cyclobutyl, cyclopropylmethyl, allyl and vinyl; wherein each of R^{12} and R^{13} is independently selected from hydrido, methyl, ethyl, n -propyl, isopropyl, cyclopropylmethyl, phenyl, benzyl, monochlorophenyl and dichlorophenyl; or a pharmaceutically-acceptable salt thereof.

19. The method of Claim 18 wherein said compound is of Formula II




wherein r is zero or two; wherein q is two or three; wherein R^2 is selected from hydrido, methyl, ethyl and phenyl; wherein R^3 is selected from hydrido, cyclohexylmethyl, benzyl, fluorobenzyl, chlorobenzyl,


fluoronaphthylmethyl and chloronaphthylmethyl; wherein each of R⁴ and R⁶ is hydrido; wherein R⁵ is selected from





5 wherein V is selected from hydrido and methyl; wherein m is one or two; wherein R⁷ is cyclohexylmethyl; wherein R⁸ is selected from n-propyl, isobutyl, cyclopropyl, cyclopropylmethyl, allyl and vinyl; wherein R¹² and R¹³ is
10 independently selected from methyl, ethyl and isopropyl; or a pharmaceutically-acceptable salt thereof.

20. The method of Claim 18 wherein said compound is selected from compounds, their tautomers and
15 pharmaceutically-acceptable salts thereof, of the group consisting of:









21. The method of Claim 20 wherein said compound is N-[1R*-[[[1S,1R*-(cyclohexylmethyl)-2S*,3R*-dihydroxy-5-methylhexyl]amino]carbonyl]-3-butynyl]- α R*-[[[2-(dimethylamino)ethyl]sulfonyl]methyl]benzenepropanamide.

22. The method of Claim 15 for treating hypertension.

10 23. The method of Claim 15 for treating glaucoma.